



Institut für Mathematik

Comparing maximal mean values on different scales


by

*T. Havenith
S. Scholtes*

Report No. **76**

2015

January 2015



Institute for Mathematics, RWTH Aachen University

**Templergraben 55, D-52062 Aachen
Germany**

COMPARING MAXIMAL MEAN VALUES ON DIFFERENT SCALES

THOMAS HAVENITH AND SEBASTIAN SCHOLTES

ABSTRACT. When computing the average speed of a car over different time periods from given GPS data, it is conventional wisdom that the maximal average speed over all time intervals of fixed length decreases if the interval length increases. However, this intuition is wrong. This can be easily seen by the example in Figure 1 for the two time periods $T < S$. We investigate this phenomenon and make rigorous in which sense this intuition is still true.




FIGURE 1. Maximal average speeds for larger time periods are not necessarily smaller.

In many applications, rates are measured as average velocities; for instance the speed monitored by Speed Check Services SPECS in the UK, capacities of rides in amusement parks and the drift velocity of electrons are obtained in this way [Smi12, Ahm97, Gri99]. In other fields mean values over different time intervals and their relation are important:

- An athlete has a high heart rate during a competition and maybe even a very high heart rate for a small time during this contest, but having a similar heart rate for a whole week would result in serious health problems.
- While approximately 10 percent of people that are exposed to a radiation dose of 1 Sv over a short time period die from light radiation poisoning in the first 30 days [Har11, p.400], this is also the maximal dose allowed for NASA astronauts over their career, [NAS08, p.7].
- Certain materials, for example a power line, can endure fairly high stresses, currents etc. for a small time interval, but the average load on a larger time scale has to be much smaller. Another illustration for the effect of the time scale is the phenomenon of creep [MC99].

All these examples show that one expects maximal mean values to become smaller as the interval size decreases. We study mean values (in the discrete as well as in the continuous setting) with regard to this property and find that it is not true in general (Lemmata 3 and 8), but that it is true if one compares the maximal mean value over all regions of size T with those taken on regions of size $d \cdot T$ for $d \in \mathbb{N}$ (Lemmata 3 and 7).

Date: January 20, 2015.

2010 Mathematics Subject Classification. 46B25; 62P30, 46A45, 47A30.

Key words and phrases. mean value, arithmetic mean, norm, L_p norm.

1. DISCRETE SETTING

Let ℓ^∞ be the space of bounded real sequences. For $x \in \ell^\infty$, $p \in (0, \infty)$ and $n \in \mathbb{N}$ we define

$$\|x\|_{\infty,p,n} := \sup_{j \in \mathbb{N}} \left(\frac{1}{n} \sum_{i=j}^{j+(n-1)} |x_i|^p \right)^{\frac{1}{p}}.$$

Lemma 1 (For $p \in [1, \infty)$ the mapping $\|\cdot\|_{\infty,p,n}$ is an equivalent norm to $\|\cdot\|_\infty$).

Let $p \in [1, \infty)$, $n \in \mathbb{N}$. Then $\|\cdot\|_{\infty,p,n}$ is a norm on ℓ^∞ with

$$n^{-\frac{1}{p}} \|x\|_\infty \leq \|x\|_{\infty,p,n} \leq \|x\|_\infty.$$

Proof. Clearly $\|\cdot\|_{\infty,p,n}$ is positively homogenous of degree one and $\|x\|_{\infty,p,n} \geq 0$ with equality if and only if $x = 0$. From the triangle inequality of the p -norm in Euclidean n -space we obtain the triangle inequality for $\|\cdot\|_{\infty,p,n}$. Additionally, we obtain

$$n^{-\frac{1}{p}} \|x\|_\infty = \left(\frac{1}{n} \|x\|_\infty^p \right)^{\frac{1}{p}} \leq \|x\|_{\infty,p,n} = \sup_{j \in \mathbb{N}} \left(\frac{1}{n} \sum_{i=j}^{j+(n-1)} |x_i|^p \right)^{\frac{1}{p}} \leq \left(\frac{1}{n} n \|x\|_\infty^p \right)^{\frac{1}{p}} = \|x\|_\infty.$$

□

This means that the spaces $(\ell^\infty, \|\cdot\|_{\infty,p,n})$ are non-reflexive Banach spaces with the same topology as the standard ℓ^∞ .

Lemma 2 (Inequality for norms).

Let $\alpha_i \in \mathbb{N}$ with $n = \sum_{l=1}^d \alpha_l$. Then for all $p \in [1, \infty)$ and $x \in \ell^\infty$ we have that

$$\|x\|_{\infty,p,n}^p \leq \sum_{l=1}^d \frac{\alpha_l}{n} \|x\|_{\infty,p,\alpha_l}^p.$$

Proof. Let $j \in \mathbb{N}$, $\varepsilon > 0$ such that $\|x\|_{\infty,p,n}^p - \varepsilon \leq \frac{1}{n} \sum_{i=j}^{j+(n-1)} x_i^p$. Then

$$\|x\|_{\infty,p,n}^p - \varepsilon \leq \frac{1}{n} \sum_{i=j}^{j+(n-1)} x_i^p = \frac{1}{n} \sum_{l=1}^d \sum_{i=j+\sum_{k=1}^{l-1} \alpha_k}^{j+\sum_{k=1}^l \alpha_k + (\alpha_l-1)} x_i^p \leq \frac{1}{n} \sum_{l=1}^d \alpha_l \|x\|_{\infty,p,\alpha_l}^p.$$

□

Lemma 3 (Norms $\|\cdot\|_{\infty,p,dn}$ of multiples of n are ordered).

Let $m, n \in \mathbb{N}$ and $n \leq m$.

- If n is a factor of m , i.e. $m = dn$ for $d \in \mathbb{N}$ then

$$\|x\|_{\infty,p,dn} = \|x\|_{\infty,p,m} \leq \|x\|_{\infty,p,n} \quad \text{for all } x \in \ell^\infty.$$

- If n is no factor of m then there are $x, y \in \ell^\infty$ such that

$$\|x\|_{\infty,p,n} < \|x\|_{\infty,p,m} \quad \text{and} \quad \|y\|_{\infty,p,m} < \|y\|_{\infty,p,n}.$$

Proof. If we choose all $\alpha_l = n$ in Lemma 2 we immediately obtain the first item. For the second item define $x(n) \in \ell^\infty$ by $x(n)_i = 1$ if $i \in \{(d-1)n+1 \mid d \in \mathbb{N}\}$ and $x(n)_i = 0$ otherwise. Then

$$\|x(n)\|_{\infty,p,n}^p = \frac{1}{n} < \frac{d}{dn-1} \leq \frac{d}{(d-1)n+j} = \|x(n)\|_{\infty,p,(d-1)n+j}^p$$

for $j = 1, \dots, n-1$ and $d \in \mathbb{N}$. Note that the natural numbers of the form $m = (d-1)n+j$ for $j = 1, \dots, n-1$, $d \in \mathbb{N}$ are exactly those which are either smaller than n or such that n is no factor of m . This proves the proposition as we obtain the opposite inequality by interchanging the roles of m and n . □

In the following Lemma we show that counterexamples, however, cannot be too bad.

Lemma 4 (Counterexamples cannot be too bad).

Let $p \in [1, \infty)$, $n < m$. Then for all $x \in \ell^\infty$ we have that

$$\|x\|_{\infty, p, m}^p \leq \frac{(\lfloor \frac{m}{n} \rfloor + 1)n}{m} \|x\|_{\infty, p, n}^p \leq 2 \|x\|_{\infty, p, n}^p.$$

Proof. Choose $d := \lfloor \frac{m}{n} \rfloor$ and $\varepsilon > 0$. Then there is $j \in \mathbb{N}$ such that

$$\begin{aligned} \|x\|_{\infty, p, m}^p - \varepsilon &\leq \frac{1}{m} \sum_{i=j}^{j+(m-1)} |x_i|^p \leq \frac{1}{m} \sum_{i=j}^{j+((d+1)n-1)} |x_i|^p \\ &\leq \frac{(d+1)n}{m} \|x\|_{\infty, p, (d+1)n}^p \leq \frac{(d+1)n}{m} \|x\|_{\infty, p, n}^p \end{aligned}$$

according to Lemma 3. \square

2. CONTINUOUS SETTING

Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be a function. For $V > 0$, $p \in (0, \infty)$ we define

$$\|f\|_{p, V} := \sup_{\substack{\Omega \subset \mathbb{R}^N \text{ open} \\ \Omega \text{ convex} \\ |\Omega|=V}} \left(\frac{1}{|\Omega|} \int_{\Omega} |f(x)|^p dx \right)^{\frac{1}{p}}.$$

Note that this norm bears a resemblance to the Hardy-Littlewood maximal function from harmonic analysis. If we denote the space of all functions $f : \mathbb{R}^N \rightarrow \mathbb{R}$ with $f = 0$ a.e. by \mathcal{N} we can define

$$L_V^p := \{f : \mathbb{R}^N \rightarrow \mathbb{R} \mid \|f\|_{p, V} < \infty\}/\mathcal{N}.$$

By a slight abuse of notation we also want to consider $\|\cdot\|_{p, V}$ as a mapping from L_V^p to \mathbb{R} in the obvious manner.

Lemma 5 (For $p \in [1, \infty)$ the mapping $\|\cdot\|_{p, V}$ is a norm on L_V^p).

Let $p \in [1, \infty)$, $V > 0$. Then $\|\cdot\|_{p, V}$ is a norm on L_V^p .

Proof. Clearly $\|\cdot\|_{p, V}$ is positively homogenous of degree one and $\|f\|_{p, V} \geq 0$ with equality if and only if $f = 0$ a.e.. Now, the triangle inequality for L^p proves the triangle inequality for $\|\cdot\|_{p, V}$. \square

Lemma 6 (Inequality for norms).

Let $\alpha_i > 0$ with $V = \sum_{l=1}^d \alpha_l$. Then for all $p \in [1, \infty)$, $V > 0$ and $f \in L_V^p$ holds

$$\|f\|_{p, V}^p \leq \sum_{l=1}^d \frac{\alpha_l}{V} \|f\|_{p, \alpha_l}^p.$$

Proof. Step 1 Let $\varepsilon > 0$ and $\Omega \subset \mathbb{R}^N$ be open and connected such that $\|f\|_{p, V}^p - \varepsilon \leq \frac{1}{|\Omega|} \int_{\Omega} |f(x)|^p dx$. According to Step 2 there is a hyperplane that cuts Ω into two convex sets Ω_1 and $\tilde{\Omega}_1$ with $|\Omega_1| = \alpha_1$ and $|\tilde{\Omega}_1| = \sum_{l=2}^d \alpha_l$. Now iterating this procedure with $\tilde{\Omega}_1$ we finally obtain d open convex sets Ω_i with $|\Omega_i| = \alpha_i$ and $\Omega = \bigcup_{l=1}^d \Omega_i \cup A$, where $\mathcal{H}^N(A) = 0$, as A is contained in a finite union of hyperplanes. Therefore

$$\|f\|_{p, V}^p - \varepsilon \leq \frac{1}{|\Omega|} \int_{\Omega} |f(x)|^p dx = \frac{1}{|\Omega|} \sum_{l=1}^d \int_{\Omega_i} |f(x)|^p dx \leq \frac{1}{|\Omega|} \sum_{l=1}^d \alpha_l \|f\|_{p, \alpha_l}^p.$$

Step 2 Let Ω be an open convex set and $\alpha + \beta = |\Omega|$ for $\alpha, \beta > 0$. If we define

$$\Omega_s := \{x \in \Omega \mid x_1 = s\}$$

we know by Cavalieri's Principle (a special case of Fubini's Theorem) that the function $\varphi(t) := \int_{-\infty}^t |\Omega_s| ds$ measures the volume of the set Ω intersected with the open half-space below (in e_1 direction) the hyperplane $\{x \in \mathbb{R}^N \mid x_1 = t\}$. Additionally $\lim_{t \rightarrow -\infty} \varphi(t) = 0$, $\lim_{t \rightarrow \infty} \varphi(t) = |\Omega|$ and φ is continuous. Therefore the intermediate value theorem yields the existence of a t_0 such that $\Omega_1 = \bigcup_{t \in (-\infty, t_0)} \Omega_t$ and $\tilde{\Omega}_1 = \bigcup_{t \in (t_0, \infty)} \Omega_t$ are open convex sets (as intersection of two such sets) and $|\Omega_1| = \alpha$ and $|\tilde{\Omega}_1| = |\Omega| - \alpha = \beta$. \square

Lemma 7 (Norms $\|\cdot\|_{p,d,V}$ of multiples of V are ordered).

Let $p \in [1, \infty)$, $V > 0$ and $d \in \mathbb{N}$. Then

$$\|f\|_{p,d,V} \leq \|f\|_{p,V} \quad \text{for all } f \in L_V^p.$$

Proof. If we chose all $\alpha_l = V$ in Lemma 6 we directly obtain the result. \square

Lemma 8 (For $N = 1$, $T < S$ and T no factor of S the norms are not ordered).

Let $N = 1$, $p \in [1, \infty)$ and $T, S > 0$, $T < S$ such that T is no factor of S . Then there are $f, g \in L_T^p$ such that

$$\|f\|_{p,T} < \|f\|_{p,S} \quad \text{and} \quad \|g\|_{p,S} < \|g\|_{p,T}.$$

Proof. Let $T < S$ such that T is not a factor of S , i.e. $\frac{S}{T} \notin \mathbb{N}$. Write $d := \lfloor \frac{S}{T} \rfloor$. Then it is possible to choose $\varepsilon > 0$ such that $(d+1)\varepsilon < S - dT$. Let $\psi_\varepsilon(x)$ be a bump function of L^p mass 1 with support in $(0, \varepsilon)$. Then $f(x) = \sum_{i=0}^d \psi_\varepsilon(x - i(T + \varepsilon))$ is an L_T^p function and

$$\|f\|_{p,T}^p = \frac{1}{T} < \frac{d+1}{S} = \|f\|_{p,S}^p,$$

because $\frac{S}{T} < \lfloor \frac{S}{T} \rfloor + 1$. On the other hand, $g = \psi_\varepsilon \in L_T^p$ yields

$$\|g\|_{p,S}^p = \frac{1}{S} < \frac{1}{T} = \|g\|_{p,T}^p.$$

\square

Corollary 9 (Counterexamples cannot be too bad).

Let $p \in [1, \infty)$, $T < S$. Then for all $f \in L_S^p$ holds

$$\|f\|_{p,S}^p \leq \frac{(\lfloor \frac{S}{T} \rfloor + 1)T}{S} \|f\|_T^p \leq 2 \|f\|_T^p.$$

Proof. Choose $d := \lfloor \frac{S}{T} \rfloor$ and $\varepsilon > 0$ such that $\|f\|_{p,S}^p - \varepsilon \leq \frac{1}{S} \int_{\Omega} |f(t)|^p dt$. Then we can extend Ω convexly to an open $\tilde{\Omega}$ with $\Omega \subset \tilde{\Omega}$ and $|\tilde{\Omega}| = (d+1)T$. This can for example be done by taking the convex hull $\tilde{\Omega}_y$ of Ω and a point y moving along a normal at a fixed point $y \in \partial\Omega$. By an argument similar to that in the proof of Lemma 6 Step 2, we see that x can be chosen in such a way that $|\tilde{\Omega}_y| = (d+1)T$. Now

$$\|f\|_{p,S}^p - \varepsilon \leq \frac{1}{S} \int_{\Omega} |f(t)|^p dt \leq \frac{1}{S} \int_{\tilde{\Omega}} |f(t)|^p dt \leq \frac{(d+1)T}{S} \|f\|_{(d+1)T}^p \leq \frac{(d+1)T}{S} \|f\|_T^p$$

by Lemma 7. \square

REFERENCES

- [Ahm97] Reza H. Ahmadi, *Opportunities for simulation in amusement parks*, Oper. Res. **45** (1997), no. 1, 1–13.
- [Gri99] David Griffiths, *Introduction to electrodynamics*, Prentice-Hall, 1999.
- [Har11] Gavin D. J. Harper, *Green Health: An A-to-Z Guide*, ch. Radiation Sources, pp. 374–401, SAGE, 2011.
- [MC99] M. A. Meyers and K. K. Chawla, *Mechanical Behavior of Materials*, Cambridge University Press, 1999.
- [NAS08] NASA, *Space Faring: The Radiation Challenge Introduction and Module 1: Radiation*, Educator Guide EP-2008-08-116-MSFC, NASA, 2008.
- [Smi12] B.D.V. Smith, *Vehicle speed monitoring system*, US Patent 8,189,048, 2012.

THOMAS HAVENITH, INFRASTRUKTURPLANUNG (I.ETZ 1), DB ENERGIE GMBH, PFARRER-PERABO-PLATZ 2, 60326 FRANKFURT AM MAIN, GERMANY

E-mail address: thomas.havenith@deutschebahn.com

URL: <http://www.dbenergie.de/>

SEBASTIAN SCHOLTES, INSTITUT FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, TEMPLERGRABEN 55, 52062 AACHEN, GERMANY

E-mail address: sebastian.scholtes@rwth-aachen.de

URL: <http://www.instmath.rwth-aachen.de/~scholtes/home/>

Reports des Instituts für Mathematik der RWTH Aachen

- [1] Bemelmans J.: *Die Vorlesung "Figur und Rotation der Himmelskörper"* von F. Hausdorff, WS 1895/96, Universität Leipzig, S 20, März 2005
- [2] Wagner A.: *Optimal Shape Problems for Eigenvalues*, S 30, März 2005
- [3] Hildebrandt S. and von der Mosel H.: *Conformal representation of surfaces, and Plateau's problem for Cartan functionals*, S 43, Juli 2005
- [4] Reiter P.: *All curves in a C^1 -neighbourhood of a given embedded curve are isotopic*, S 8, Oktober 2005
- [5] Maier-Paape S., Mischaikow K. and Wanner T.: *Structure of the Attractor of the Cahn-Hilliard Equation*, S 68, Oktober 2005
- [6] Strzelecki P. and von der Mosel H.: *On rectifiable curves with L^p bounds on global curvature: Self-avoidance, regularity, and minimizing knots*, S 35, Dezember 2005
- [7] Bandle C. and Wagner A.: *Optimization problems for weighted Sobolev constants*, S 23, Dezember 2005
- [8] Bandle C. and Wagner A.: *Sobolev Constants in Disconnected Domains*, S 9, Januar 2006
- [9] McKenna P.J. and Reichel W.: *A priori bounds for semilinear equations and a new class of critical exponents for Lipschitz domains*, S 25, Mai 2006
- [10] Bandle C., Below J. v. and Reichel W.: *Positivity and anti-maximum principles for elliptic operators with mixed boundary conditions*, S 32, Mai 2006
- [11] Kyed M.: *Travelling Wave Solutions of the Heat Equation in Three Dimensional Cylinders with Non-Linear Dissipation on the Boundary*, S 24, Juli 2006
- [12] Blatt S. and Reiter P.: *Does Finite Knot Energy Lead To Differentiability?*, S 30, September 2006
- [13] Grunau H.-C., Ould Ahmedou M. and Reichel W.: *The Paneitz equation in hyperbolic space*, S 22, September 2006
- [14] Maier-Paape S., Miller U., Mischaikow K. and Wanner T.: *Rigorous Numerics for the Cahn-Hilliard Equation on the Unit Square*, S 67, Oktober 2006
- [15] von der Mosel H. and Winklmann S.: *On weakly harmonic maps from Finsler to Riemannian manifolds*, S 43, November 2006
- [16] Hildebrandt S., Maddocks J. H. and von der Mosel H.: *Obstacle problems for elastic rods*, S 21, Januar 2007
- [17] Galdi P. Giovanni: *Some Mathematical Properties of the Steady-State Navier-Stokes Problem Past a Three-Dimensional Obstacle*, S 86, Mai 2007
- [18] Winter N.: *$W^{2,p}$ and $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear, uniformly elliptic equations*, S 34, Juli 2007
- [19] Strzelecki P., Szumańska M. and von der Mosel H.: *A geometric curvature double integral of Menger type for space curves*, S 20, September 2007
- [20] Bandle C. and Wagner A.: *Optimization problems for an energy functional with mass constraint revisited*, S 20, März 2008
- [21] Reiter P., Felix D., von der Mosel H. and Alt W.: *Energetics and dynamics of global integrals modeling interaction between stiff filaments*, S 38, April 2008
- [22] Belloni M. and Wagner A.: *The ∞ Eigenvalue Problem from a Variational Point of View*, S 18, Mai 2008
- [23] Galdi P. Giovanni and Kyed M.: *Steady Flow of a Navier-Stokes Liquid Past an Elastic Body*, S 28, Mai 2008
- [24] Hildebrandt S. and von der Mosel H.: *Conformal mapping of multiply connected Riemann domains by a variational approach*, S 50, Juli 2008
- [25] Blatt S.: *On the Blow-Up Limit for the Radially Symmetric Willmore Flow*, S 23, Juli 2008
- [26] Müller F. and Schikorra A.: *Boundary regularity via Uhlenbeck-Rivièvre decomposition*, S 20, Juli 2008
- [27] Blatt S.: *A Lower Bound for the Gromov Distortion of Knotted Submanifolds*, S 26, August 2008
- [28] Blatt S.: *Chord-Arc Constants for Submanifolds of Arbitrary Codimension*, S 35, November 2008
- [29] Strzelecki P., Szumańska M. and von der Mosel H.: *Regularizing and self-avoidance effects of integral Menger curvature*, S 33, November 2008
- [30] Gerlach H. and von der Mosel H.: *Yin-Yang-Kurven lösen ein Packungsproblem*, S 4, Dezember 2008
- [31] Buttazzo G. and Wagner A.: *On some Rescaled Shape Optimization Problems*, S 17, März 2009
- [32] Gerlach H. and von der Mosel H.: *What are the longest ropes on the unit sphere?*, S 50, März 2009
- [33] Schikorra A.: *A Remark on Gauge Transformations and the Moving Frame Method*, S 17, Juni 2009
- [34] Blatt S.: *Note on Continuously Differentiable Isotopies*, S 18, August 2009
- [35] Knappmann K.: *Die zweite Gebietsvariation für die gebeulte Platte*, S 29, Oktober 2009
- [36] Strzelecki P. and von der Mosel H.: *Integral Menger curvature for surfaces*, S 64, November 2009
- [37] Maier-Paape S., Imkeller P.: *Investor Psychology Models*, S 30, November 2009
- [38] Scholtes S.: *Elastic Catenoids*, S 23, Dezember 2009
- [39] Bemelmans J., Galdi G.P. and Kyed M.: *On the Steady Motion of an Elastic Body Moving Freely in a Navier-Stokes Liquid under the Action of a Constant Body Force*, S 67, Dezember 2009
- [40] Galdi G.P. and Kyed M.: *Steady-State Navier-Stokes Flows Past a Rotating Body: Leray Solutions are Physically Reasonable*, S 25, Dezember 2009

- [41] Galdi G.P. and Kyed M.: *Steady-State Navier-Stokes Flows Around a Rotating Body: Leray Solutions are Physically Reasonable*, S 15, Dezember 2009
- [42] Bemelmans J., Galdi G.P. and Kyed M.: *Fluid Flows Around Floating Bodies, I: The Hydrostatic Case*, S 19, Dezember 2009
- [43] Schikorra A.: *Regularity of $n/2$ -harmonic maps into spheres*, S 91, März 2010
- [44] Gerlach H. and von der Mosel H.: *On sphere-filling ropes*, S 15, März 2010
- [45] Strzelecki P. and von der Mosel H.: *Tangent-point self-avoidance energies for curves*, S 23, Juni 2010
- [46] Schikorra A.: *Regularity of $n/2$ -harmonic maps into spheres (short)*, S 36, Juni 2010
- [47] Schikorra A.: *A Note on Regularity for the n -dimensional H -System assuming logarithmic higher Integrability*, S 30, Dezember 2010
- [48] Bemelmans J.: *Über die Integration der Parabel, die Entdeckung der Kegelschnitte und die Parabel als literarische Figur*, S 14, Januar 2011
- [49] Strzelecki P. and von der Mosel H.: *Tangent-point repulsive potentials for a class of non-smooth m -dimensional sets in \mathbb{R}^n . Part I: Smoothing and self-avoidance effects*, S 47, Februar 2011
- [50] Scholtes S.: *For which positive p is the integral Menger curvature \mathcal{M}_p finite for all simple polygons*, S 9, November 2011
- [51] Bemelmans J., Galdi G. P. and Kyed M.: *Fluid Flows Around Rigid Bodies, I: The Hydrostatic Case*, S 32, Dezember 2011
- [52] Scholtes S.: *Tangency properties of sets with finite geometric curvature energies*, S 39, Februar 2012
- [53] Scholtes S.: *A characterisation of inner product spaces by the maximal circumradius of spheres*, S 8, Februar 2012
- [54] Kolasiński S., Strzelecki P. and von der Mosel H.: *Characterizing $W^{2,p}$ submanifolds by p -integrability of global curvatures*, S 44, März 2012
- [55] Bemelmans J., Galdi G.P. and Kyed M.: *On the Steady Motion of a Coupled System Solid-Liquid*, S 95, April 2012
- [56] Deipenbrock M.: *On the existence of a drag minimizing shape in an incompressible fluid*, S 23, Mai 2012
- [57] Strzelecki P., Szumańska M. and von der Mosel H.: *On some knot energies involving Menger curvature*, S 30, September 2012
- [58] Overath P. and von der Mosel H.: *Plateau's problem in Finsler 3-space*, S 42, September 2012
- [59] Strzelecki P. and von der Mosel H.: *Menger curvature as a knot energy*, S 41, Januar 2013
- [60] Strzelecki P. and von der Mosel H.: *How averaged Menger curvatures control regularity and topology of curves and surfaces*, S 13, Februar 2013
- [61] Hafizogullari Y., Maier-Paape S. and Platen A.: *Empirical Study of the 1-2-3 Trend Indicator*, S 25, April 2013
- [62] Scholtes S.: *On hypersurfaces of positive reach, alternating Steiner formulæ and Hadwiger's Problem*, S 22, April 2013
- [63] Bemelmans J., Galdi G.P. and Kyed M.: *Capillary surfaces and floating bodies*, S 16, Mai 2013
- [64] Bandle, C. and Wagner A.: *Domain derivatives for energy functionals with boundary integrals; optimality and monotonicity*, S 13, Mai 2013
- [65] Bandle, C. and Wagner A.: *Second variation of domain functionals and applications to problems with Robin boundary conditions*, S 33, Mai 2013
- [66] Maier-Paape, S.: *Optimal f and diversification*, S 7, Oktober 2013
- [67] Maier-Paape, S.: *Existence theorems for optimal fractional trading*, S 9, Oktober 2013
- [68] Scholtes, S.: *Discrete Möbius Energy*, S 11, November 2013
- [69] Bemelmans, J.: *Optimale Kurven – über die Anfänge der Variationsrechnung*, S 22, Dezember 2013
- [70] Scholtes, S.: *Discrete Thickness*, S 12, Februar 2014
- [71] Bandle, C. and Wagner A.: *Isoperimetric inequalities for the principal eigenvalue of a membrane and the energy of problems with Robin boundary conditions.*, S 12, März 2014
- [72] Overath P. and von der Mosel H.: *On minimal immersions in Finsler space.*, S 26, April 2014
- [73] Bandle, C. and Wagner A.: *Two Robin boundary value problems with opposite sign.*, S 17, Juni 2014
- [74] Knappmann, K. and Wagner A.: *Optimality conditions for the buckling of a clamped plate.*, S 23, September 2014
- [75] Bemelmans, J.: *Über den Einfluß der mathematischen Beschreibung physikalischer Phänomene auf die Reine Mathematik und die These von Wigner*, S 23, September 2014
- [76] Harenith, T. and Scholtes, S.: *Comparing maximal mean values on different scales*, S 4, Januar 2015