

Institut für Mathematik

Two Robin boundary value problems with opposite
sign.


by

C. Bandle
A. Wagner

Report No. **73**

2014

June 2014



Institute for Mathematics, RWTH Aachen University

Templergraben 55, D-52062 Aachen
Germany

Two Robin boundary value problems with opposite sign.

June 12, 2014

CATHERINE BUNDLE

Mathematisches Institut, Universität Basel,
Rheinsprung 21, CH-4051 Basel, Switzerland

ALFRED WAGNER

Institut für Mathematik, RWTH Aachen
Templergraben 55, D-52062 Aachen, Germany

Abstract

Key words: Local Rayleigh-Faber Krahn inequalities, Robin boundary conditions, domain variation, harmonic transplantation.

MSC2010: 49K20, 49R05, 15A42, 35J20, 35N25.

Abstract

In this paper we discuss a boundary value and an eigenvalue problem with Robin boundary conditions of opposite sign. The eigenvalue problem arises in the study of the reaction-diffusion equation with dynamical boundary conditions. The dependence of the energy and the principle eigenvalues of the domains is investigated. The first and second domain variations are derived for nearly spherical domains. It is shown that in contrast to the classical Robin conditions the second variation changes sign and has singularities which depend on the eigenvalues of a Steklov problem. By means of the harmonic transplantation isoperimetric inequalities are derived for the principal eigenvalues in arbitrary domains.

1 Introduction

In this paper we discuss an elliptic boundary value problem with Robin boundary conditions of the opposite sign and a corresponding eigenvalue problem with the eigenvalue in the domain and on the boundary. The motivation comes from a classical model for the heat distribution in a body $\Omega \subset \mathbb{R}^3$, expressed by the heat equation

$$\partial_t T(x, t) - \Delta T(x, t) = 1 \text{ in } \Omega \times \mathbb{R}^+,$$

where ∂_t denotes partial differentiation with respect to t , Δ is the Laplace operator in \mathbb{R}^3 and the right-hand side describes a source of constant density 1. We suppose that the body is immersed into an ice bath and that on the boundary there is a regulating system with the following property: if the temperature drops below zero heat is carried into and otherwise out of the body, according to the law $\partial_\nu T = -\alpha T$ on $\partial\Omega$. Here ∂_ν denotes the outer normal derivative of Ω and $\alpha \in \mathbb{R}^+$ is a fixed positive number. This boundary condition is of the *Neumann-Robin* type. Observe that this flux condition is different from the classical Newton's law where α is of opposite sign.

We shall also consider the case where at the boundary a reaction takes place. Then on $\partial\Omega$, T satisfies the *dynamical boundary conditions* $\partial_\nu T = -\sigma \partial_t T$ where $\sigma < 0$.

In the first case the stationary solutions satisfy the elliptic boundary value problem

$$(1.1) \quad \Delta u + 1 = 0 \text{ in } \Omega, \quad \partial_\nu u = \alpha u \text{ on } \partial\Omega.$$

Under suitable regularity assumptions it has a unique solution provided α does not coincide with a eigenvalue $0 = \mu_1 < \mu_2 \leq \dots$ of the *Steklov problem*

$$(1.2) \quad \Delta \phi = 0 \text{ in } \Omega, \quad \partial_\nu \phi = \mu \phi \text{ on } \partial\Omega.$$

If $\alpha = \mu_i$ then the solutions are not unique. To (1.1) we associate the energy

$$\mathcal{E}(\Omega) := E(u, \Omega) \text{ where } E(\Omega, v) = \int_{\Omega} |\nabla v|^2 dx - 2 \int_{\Omega} v dx - \alpha \int_{\partial\Omega} v^2 dS.$$

It is well-known that the solution of (1.1) is a critical point of $E(v, \Omega)$ in $W^{1,2}(\Omega)$ in the sense that the Frechet derivative vanishes. However it is not a local extremum.

In this paper we study the dependence of $\mathcal{E}(\Omega)$ on the domain Ω . In contrast to the case $\alpha < 0$ the techniques based on the minimum principle $\mathcal{E}(\Omega) = \min_{W^{1,2}(\Omega)} E(v, \Omega)$ do not apply. Therefore, in the spirit of the previous investigations in [5] and [6], we compute the first and second order shape derivative and discuss the behavior under small perturbations of the domain which are volume preserving. This techniques fails if α is a Steklov eigenvalue. It turns out that the first variation vanishes if Ω is a ball. The second variation for the ball depends on $\oint_{\partial B_R} (v \cdot \nu)^2 dS$ and on α . It is singular if $\alpha R = \mu_k$ and if the k th-Fourier coefficient of $(v \cdot \nu)$ with respect to orthonormal system of Steklov eigenfunctions does not vanish. At this point the second variation changes sign. This is in contrast to the case $\alpha < 0$ discussed in [5] where the ball is a local minimum for all $\alpha < 0$. In a recent paper Bucur and Giacomini [7] have shown that it is also a global minimum for the same choice of α . This first section completes the investigations in our previous paper [6] where the case $\alpha < 0$ is discussed.

The eigenvalue problem we are dealing with has its origin in the heat equation with dynamical boundary conditions. A standard method to find solutions is the

separation of variables. In fact $T(x, t) = e^{-\lambda t} \phi(x)$ solves the homogeneous problem if ϕ is an eigenfunction of

$$(1.3) \quad \Delta \phi + \lambda \phi = 0 \text{ in } \Omega, \quad \partial_\nu \phi = \sigma \lambda \phi \text{ on } \partial \Omega.$$

It was shown in [4] that there exist two sequences of eigenvalues

$$\dots \lambda_{-k} \leq \dots \lambda_{-2} \leq \lambda_{-1} < 0 = \lambda_0 < \lambda_1 \leq \lambda_2 \dots,$$

such that $\lim_{n \rightarrow \infty} \lambda_{-n} = -\infty$ and $\lim_{n \rightarrow \infty} \lambda_n = \infty$. These eigenvalues can be characterized by a min-max principle, cf. [4].

In the second part of this paper we study the domain dependence of λ_1 and λ_{-1} . By means of the harmonic transplantation [8] a global upper bound for $\lambda_1(\Omega)$ and a global lower bound for λ_{-1} is derived. They are expressed in terms of the *harmonic radius* which plays an essential role. Both bounds are sharp. In [3] it was shown that for small σ the ball has for all domains of given volume the smallest λ_1 . The situation is more involved if λ_{-1} . An investigation is carried out for nearly spherical domains of prescribed volume by means of the first and second domain variations.

Throughout this paper we shall assume that Ω is a bounded Lipschitz domain. This guarantees that the embedding $W^{1,2}(\Omega)$ into $L^2(\Omega)$ as well as the trace operator $\Gamma : W^{1,2}(\Omega) \rightarrow L^2(\partial \Omega)$ is compact. Under this condition both problems (1.1) and (1.3) are solvable in $W^{1,2}(\Omega)$ in the classical sense.

2 The boundary value problem

2.1 First domain variation

Consider problem (1.1) in a class of domains Ω_t which are small perturbations of Ω . We assume that

$$(2.1) \quad \bar{\Omega}_t = \left\{ y : y = x + tv(x) + \frac{t^2}{2}w(x) + o(t^2) : x \in \bar{\Omega} \right\}.$$

where v and w are vector fields such that

$$v, w : \bar{\Omega} \rightarrow \mathbb{R}^n \quad \text{are in } C^1(\bar{\Omega}).$$

The solutions of (1.1) in Ω_t will be denoted by $u(y; t)$ and the corresponding energy by $\mathcal{E}(t)$.

Example 1 *If Ω is the ball B_R of radius R then for all α there exists a unique radial solution*

$$u(r) = \frac{R^2}{2n} - \frac{R}{\alpha n} - \frac{r^2}{2n}.$$

The corresponding energy is

$$\mathcal{E}(B_R) = - \int_{B_R} u \, dx = |B_R| \left(\frac{R}{\alpha n} - \frac{R^2}{n(n+2)} \right).$$

If α is not a Steklov eigenvalue $u(y, ; t)$ is continuous and continuously differentiable in t . Under this condition it was shown in [5] that the first domain variation is

$$\dot{\mathcal{E}}(0) = \int_{\partial\Omega} (v \cdot \nu)[|\nabla u|^2 - 2u - 2\alpha^2 u^2 - \alpha(n-1)u^2 H] dS,$$

where H is the mean curvature of $\partial\Omega$ and u is the solution of (1.1) in Ω .

Example 2 If $\Omega = B_R$ then

$$(2.2) \quad \dot{\mathcal{E}}(0) = \left(\frac{(n+1)R}{\alpha n^2} - \frac{R^2}{n^2} \right) \int_{\partial B_R} (v \cdot \nu) dS.$$

This leads to the following

Corollary 1 Let Ω_t be a family of nearly spherical domains with prescribed volume $|\Omega_t| = |B_R|$. Then $\dot{\mathcal{E}}(0) = 0$.

Proof The volume of Ω_t is given by $\int_{B_R} J(t) dx$ where $J(t)$ is the Jacobian determinant corresponding to the transformation $y : B_R \rightarrow \Omega_t$. The Jacobian matrix corresponding to this transformation is up to second order terms of the form

$$I + tD_v + \frac{t^2}{2}D_w, \text{ where } (D_v)_{ij} = \partial_j v_i \text{ and } \partial_j = \partial/\partial x_j.$$

By Jacobi's formula we have for small t

$$(2.3) \quad \begin{aligned} J(t) &:= \det \left(I + tD_v + \frac{t^2}{2}D_w \right) \\ &= 1 + t \operatorname{div} v + \frac{t^2}{2} \left((\operatorname{div} v)^2 - D_v : D_v + \operatorname{div} w \right) + o(t^2). \end{aligned}$$

Here we used the notation

$$D_v : D_v := \partial_i v_j \partial_j v_i.$$

Hence

$$|\Omega_t| = \int_{B_R} J(t) dx = |B_R| + t \int_{B_R} \operatorname{div} v dx + \frac{t^2}{2} \int_{B_R} \left((\operatorname{div} v)^2 - D_v : D_v + \operatorname{div} w \right) dx + o(t^2).$$

For the first variation we have only to require that y is volume preserving of the first order, that is

$$(2.4) \quad \int_{B_R} \operatorname{div} v dx = \int_{\partial B_R} (v \cdot \nu) dS = 0.$$

This together with (2.2) proves the assertion. \square

A further consequence of (2.2) is the local monotonicity property

Corollary 2 If $0 < \alpha R < n + 1$ and $|\Omega_t| > |B_R|$ then $\dot{\mathcal{E}}(0) > 0$, otherwise if $\alpha R > n + 1$ then $\dot{\mathcal{E}}(0) < 0$.

Proof By our assumption we have $\int_{\partial B_R} (v \cdot \nu) dS > 0$ The sign of $\dot{\mathcal{E}}(0)$ depends therefore on the sign of $(n+1)\alpha R - (\alpha R)^2$. \square

2.2 Second domain variation for nearly spherical domains

Corollary 1 gives rise to the following question: *is $\mathcal{E}(B_R)$ a local extremum among the family $(\Omega_t)_t$ of perturbed domains with the same volume as B_R ?* The answer will be obtained from the second variation.

Let $u_t(x) := u(y(x); t)$ be the solution of $\Delta u + 1 = 0$ in Ω_t , $\partial_\nu u = \alpha u$ on $\partial\Omega_t$ transformed onto Ω . If u_t is differentiable then

$$\frac{d}{dt}u_t(x)|_{t=0} = u'(x) + v \cdot \nabla u,$$

where $u = u_0$ is the solution of (1.1) in B_R .

It was shown in [5] that formally u' solves the inhomogeneous boundary value problem

$$(2.5) \quad \Delta u' = 0 \quad \text{in } B_R$$

$$(2.6) \quad \partial_\nu u' = \alpha u' + \left(\frac{1 - \alpha R}{n} \right) v \cdot \nu \quad \text{on } \partial B_R.$$

Let us assume for the moment that such a u' exists. This is certainly the case if α does not coincide with a Steklov eigenvalue μ_i .

For the next result we consider perturbations which, in addition to the condition (2.4), satisfy the volume preservation of the second order, namely

$$(2.7) \quad \int_{B_R} ((\operatorname{div} v)^2 - D_v : D_v + \operatorname{div} w) dx = 0.$$

Set

$$Q(u') := \int_{B_R} |\nabla u'|^2 dx - \alpha \int_{\partial B_R} u'^2 dS.$$

Then the following formula was derived in [5].

Lemma 1 *Assume $\alpha \neq \mu_i$, (2.4) and (2.7). Put $\mathcal{S}(t) := |\partial\Omega_t|$. Then*

$$(2.8) \quad \ddot{\mathcal{S}}(0) = -2Q(u') + \frac{2R}{n^2}(1 - \alpha R) \int_{\partial B_R} (v \cdot \nu)^2 dS - \frac{R^2}{\alpha n^2} \ddot{\mathcal{S}}(0).$$

For a ball and for volume preserving perturbations the second variation of the surface is of the form

$$\ddot{\mathcal{S}}(0) = \oint_{\partial B_R} \left(|\nabla^\tau (v \cdot \nu)|^2 - \frac{(n-1)}{R^2} (v \cdot \nu)^2 \right) dS,$$

where ∇^τ stands for the tangential gradient on ∂B_R .

2.2.1 Discussion of $\ddot{\mathcal{E}}(0)$

We write for short

$$(2.9) \quad \mathcal{F} := -2Q_g(u') + \frac{2R}{n^2}(1 - \alpha R) \int_{\partial B_R} (v \cdot \nu)^2 dS.$$

In order to estimate \mathcal{F} we consider the Steklov eigenvalue problem (1.2). An elementary computation yields $\mu_1 = 0$, and $\mu_k = \frac{k-1}{R}$ (for $k \geq 2$ and counted without multiplicity). The second eigenvalue $\mu_2 = 1/R$ has multiplicity n and its eigenfunctions are $\frac{x_1}{R}, \dots, \frac{x_n}{R}$.

From now on we shall count the eigenvalues μ_i with their multiplicity, i.e. $\mu_2 = \mu_3 = \dots = \mu_{n+1} = 1/R$ and $\mu_{n+2} = 2/R$ etc. There exists a complete system of eigenfunctions $\{\phi_i\}_{i \geq 1}$ such that

$$(2.10) \quad \oint_{\partial B_R} \phi_i \phi_j dS = \delta_{ij}.$$

In view of the completeness we can write

$$u'(x) = \sum_{i=1}^{\infty} c_i \phi_i \quad \text{and} \quad (v \cdot \nu) = \sum_{i=1}^{\infty} b_i \phi_i.$$

Note that the first eigenfunction $\phi_1 \equiv \text{constant}$. The condition

$$0 = \oint_{\partial B_R} (v \cdot \nu) dS = \oint_{\partial B_R} \phi_1 (v \cdot \nu) dS$$

implies that $b_1 = 0$. From (2.6) we have also $c_1 = 0$. The coefficients b_i for $i \geq 2$ are determined from the boundary value problem (2.5), (2.6). In fact

$$(2.11) \quad b_i = \frac{n c_i (\mu_i - \alpha)}{1 - \alpha R} \quad \text{for } i = 2, 3, \dots$$

From the orthonormality conditions of the eigenfunctions it follows that

$$Q(u') = \sum_{i=2}^{\infty} c_i^2 (\mu_i - \alpha).$$

Inserting this into (2.9) we get

$$(2.12) \quad \mathcal{F} = 2 \sum_2^{\infty} c_i^2 (\mu_i - \alpha)^2 \left[\frac{R}{1 - \alpha R} - \frac{1}{\mu_i - \alpha} \right].$$

DISCUSSION OF $\ddot{\mathcal{S}}(0)$. Observe that

$$\mathcal{R}[\chi] = \frac{\oint_{\partial B_R} |\nabla^\tau \chi|^2 dS}{\oint_{\partial B_R} \chi^2 dS}$$

is the Rayleigh quotient of the Laplace-Beltrami operator on the $(n-1)$ -dimensional sphere of radius R . Its eigenvalues are $k(n-2+k)/R^2$, $k \in \mathbb{N}^+$. For volume

preserving perturbations of the first order ($v \cdot \nu$) is orthogonal to the first eigenfunction ($\oint_{\partial B_R} (v \cdot \nu) dS = 0$) and thus

$$\mathcal{R}[(v \cdot \nu)] \geq \frac{n-1}{R^2}.$$

Equality holds if and only if $(v \cdot \nu)$ belongs to the eigenspace spanned by $\{x_i\}_{i=1}^n$, $|x_i| = R$. This does not occur if we exclude small translations. Consequently $\tilde{S}(0) > 0$.

SIGN OF $\tilde{\mathcal{E}}(0)$ Let us write for short

$$A := \frac{2(1-\alpha R)^2}{n^2} \text{ and } d_k := \frac{R}{1-\alpha R} - \frac{R}{k-1-\alpha R} = \frac{k-2}{(1-\alpha R)(k-1-\alpha R)}.$$

Clearly

$$d_k \begin{cases} > 0 & \text{if } \alpha R < 1 \text{ or } \alpha R > k-1, \\ < 0 & \text{if } 1 < \alpha R < k-1. \end{cases}$$

By (2.11)

$$\mathcal{F} = \frac{2(1-\alpha R)^2}{n^2} \sum_2^\infty b_i^2 \left[\frac{R}{1-\alpha R} - \frac{1}{\mu_i - \alpha} \right] = A \sum_3^\infty \tilde{b}_k^2 d_k.$$

Remark 1 *The last equality is read in the following way. Assume for some $k \geq 2$ there are m identical eigenvalues*

$$\mu_k = \mu_{k_1} = \dots = \mu_{k_m}$$

The sum to the left thus contains the finite sum

$$\sum_{j=1}^m b_{k_j}^2 \left[\frac{R}{1-\alpha R} - \frac{1}{\mu_k - \alpha} \right].$$

The sum on the right side abbreviates this by setting $\tilde{b}_k^2 = \sum_{j=1}^m b_{k_j}^2$.

Let μ be the largest eigenvalue such that $\mu_\alpha := \max\{\mu_i : \mu_i < \alpha\}$. Suppose that $0 < p-1 < \alpha R < p$. Then $d_k > 0$ for $k = 1, \dots, p$ and $d_k < 0$ for $k > p$. Then \mathcal{F} can be split into a positive and negative part,

$$\mathcal{F} = \mathcal{F}^+ + \mathcal{F}^-, \text{ where } \mathcal{F}^+ = A \sum_3^p \tilde{b}_k^2 d_k \geq 0, \quad \mathcal{F}^- = A \sum_{p+1}^\infty \tilde{b}_k^2 d_k \leq 0.$$

If $0 < \alpha R < 1$ then $d_k > 0$ for all $k = 3, \dots$. Hence

$$(2.13) \quad \mathcal{F} = \mathcal{F}^+ = A \sum_3^\infty \tilde{b}_k^2 d_k.$$

Moreover

$$(2.14) \quad \begin{aligned} \mathcal{F}^+ &\leq A d_p \sum_3^p \tilde{b}_k^2 \leq A d_p \int_{\partial B_R} (v \cdot \nu)^2 dS \\ \mathcal{F}^- &\geq A d_{p+1} \sum_{p+1}^{\infty} \tilde{b}_k^2 \geq A d_{p+1} \int_{\partial B_R} (v \cdot \nu)^2 dS, \\ \mathcal{F} &\leq A \left[\frac{R}{1 - \alpha R} - \frac{1}{\mu_\alpha - \alpha} \right] \int_{\partial B_R} (v \cdot \nu)^2 dS. \end{aligned}$$

Recall that

$$\ddot{\mathcal{E}}(0) = \mathcal{F}^+ + \mathcal{F}^- - \frac{R^2}{\alpha n^2} \ddot{\mathcal{S}}(0).$$

Theorem 1 *Suppose that the perturbations y are different from translations and rotations. Under the assumptions of Lemma 1 and if $0 < \alpha R < \epsilon$ for ϵ sufficiently small then $\ddot{\mathcal{E}}(0) < 0$. The ball is a local maximizer for $\mathcal{E}(t)$.*

Proof By (2.13), \mathcal{F}^+ is bounded from above for $\alpha R < \epsilon$. Hence the last term dominates and $\ddot{\mathcal{S}}(0) < 0$. \square

Sign changes occur if α is in a neighborhood of μ_α . More precisely we have

Theorem 2 *Let $\tilde{b}_p \neq 0$. Suppose that α is close to $\mu_\alpha = p - 1$. Then there exists $\epsilon_0 > 0$ sufficiently small and depending only on v and w such that $\ddot{\mathcal{E}}(0) > 0$ if $p - 1 < \alpha < p - 1 + \epsilon_0$. On the other hand if $\tilde{b}_{p+1} \neq 0$ there exists $\epsilon_1 > 0$ sufficiently small such that $\ddot{\mathcal{E}}(0) < 0$ if $p - \epsilon_1 < \alpha < p$. If $\tilde{b}_p = 0$ in the first case then $\ddot{\mathcal{E}}(0)$ can be positive or negative depending on $(v \cdot \nu)$. The same is true in the second case if $\tilde{b}_{p+1} = 0$.*

Proof The expressions containing d_p in the first case and, d_{p+1} in the second case dominate. \square

In order to get a sharper upper bound for $\ddot{\mathcal{E}}(0)$ in terms of v we impose the "barycenter" condition

$$(2.15) \quad \oint_{\partial B_R} x (v(x) \cdot \nu(x)) dS = 0.$$

Setting $N = (v \cdot \nu)$ we get

$$\oint_{\partial B_R} |\nabla^\tau N|^2 dS \geq \frac{2n}{R^2} \oint_{\partial B_R} N^2 dS.$$

$$\text{Thus } \ddot{\mathcal{S}}(0) \geq \frac{n+1}{R^2} \oint_{\partial B_R} N^2 dS \geq \frac{n+1}{R^2} \sum_3^p \tilde{b}_k^2.$$

We then get the following upper bound for $\ddot{\mathcal{E}}(0)$.

$$(2.16) \quad \ddot{\mathcal{E}}(0) \leq \left\{ -\frac{n+1}{\alpha n^2} + 2 \frac{R(1 - \alpha R)}{n^2} - \frac{2(1 - \alpha R)^2}{n^2(\mu_\alpha - \alpha)} \right\} \sum_3^p \tilde{b}_k^2.$$

Remark 2 Observe that since (2.15) is assumed, $b_i = 0$ for $i = 1, \dots, n$. By (2.11) this implies $c_i = 0$ for $i = 1, \dots, n$. If

$$p - 1 < \alpha R < p$$

then

$$\mu_\alpha = \frac{p-1}{R}.$$

Consequently

$$(2.17) \quad \ddot{\mathcal{E}}(0) \leq -\frac{1}{\alpha n^2} \left\{ n + 1 - \frac{2\alpha R(1-\alpha R)(p-2)}{p-1-\alpha R} \right\} \sum_3^p \tilde{b}_k^2.$$

If $p = 2$ then it follows immediately that $\ddot{\mathcal{E}}(0) \leq 0$. Let $p > 2$. The second term in the brackets is monotone decreasing in αR . Thus

$$\inf_{\alpha R \in (p-1, p)} \frac{2\alpha R(1-\alpha R)(p-2)}{p-1-\alpha R} = 2p(p^2 - 3p + 2) \geq 12.$$

Hence if $n \geq 12$, $\ddot{\mathcal{E}}(0) < 0$ for αR close to p .

Example 3 Let $\Omega_t \subset \mathbb{R}^2$ be the ellipse whose boundary $\partial\Omega_t$ is given by

$$\left\{ \frac{R \cos(\theta)}{1+t}, (1+t)R \sin(\theta) \right\},$$

where (r, θ) are the polar coordinates in the plane. This ellipse has the same area as the circle B_R and can be interpreted as a perturbation described in (2.1). We have $y = x + t(-x_1, x_2) + \frac{t^2}{2}(x_1, 0) + o(t^2)$. The eigenvalues and eigenfunctions of the Steklov eigenvalue problem (1.3) are

$$\mu = \frac{m}{R} + \alpha \text{ and } \phi = r^m (a_1 \cos(m\theta) + a_2 \sin(m\theta)).$$

We have

$$(v \cdot \nu) = -R \cos(2\theta) = b_3 \phi_3,$$

and

$$\ddot{S}(0) = \oint_{\partial B_R} \left(|\nabla^\tau N|^2 - \frac{N^2}{R^2} \right) dS = 3\pi R.$$

A straightforward computation yields

$$\ddot{\mathcal{E}}(0) = \left[-\frac{3}{4\alpha} + \frac{R(1-\alpha R)}{2(2-\alpha R)} \right] \oint_{\partial B_R} (v \cdot \nu)^2 dS.$$

with $\oint_{\partial B_R} (v \cdot \nu)^2 dS = \pi R^3$. From this expression it follows immediately that

$$\ddot{\mathcal{E}}(0) \begin{cases} > 0 \text{ if } \alpha R > 2 \\ < 0 \text{ if } \alpha R < 2. \end{cases}$$

In this example there is only one coefficient \tilde{b}_3 which does not vanish. In accordance with Theorem 2, $\ddot{\mathcal{E}}(0)$ has only one singularity at $\alpha = \mu_3 = \frac{2}{R}$.

3 Eigenvalue problem

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary. We consider the eigenvalue problem

$$(3.1) \quad \Delta u + \lambda u = 0 \quad \text{in } \Omega, \quad \partial_\nu u = \lambda \sigma u \quad \text{in } \partial\Omega.$$

σ is a negative number. Clearly $\lambda_0 = 0$ is a eigenvalue and the corresponding eigenfunction is $u_0 = \text{const.}$. We define

$$\sigma_0(\Omega) := -\frac{|\Omega|}{|\partial\Omega|}.$$

In [4] [3] the following properties were shown.

- (P1) There exists an infinite sequence of positive eigenvalues $(\lambda_n)_n$ with $\lim_{n \rightarrow \infty} \lambda_n = \infty$.
- (P2) There exists an infinite sequence of negative eigenvalues $(\lambda_{-n})_n$ with $\lim_{n \rightarrow \infty} \lambda_{-n} = -\infty$.
- (P3) If $\sigma < \sigma_0(\Omega)$, then λ_1 is simple and the corresponding eigenfunction u_1 is of constant sign.
- (P4) If $0 > \sigma > \sigma_0(\Omega)$, then λ_{-1} is simple and the corresponding eigenfunction u_{-1} is of constant sign.
- (P5) Let B_R be a ball such that $|B_R| = |\Omega|$. If $\sigma < \sigma_0(B_R)$, then $\lambda_1(\Omega) \geq \lambda_1(B_R)$.
- (P6) For any domain Ω (with $|B_R| = |\Omega|$), there exist a number $\hat{\sigma} \in (\sigma_0(\Omega), 0)$ such that $\lambda_{-1}(\Omega) \geq \lambda_{-1}(B_R)$ whenever $\sigma \in (\sigma_0(\Omega), \hat{\sigma})$.

Remark 3 For (P5) we note that the condition $\sigma < \sigma_0(B_R)$ is more restrictive than the condition $\sigma < \sigma_0(\Omega)$ if $|\Omega| = |B_R|$. This is a consequence of the isoperimetric inequality.

From (3.1) we obtain a representation formula for λ_{-1} and λ_1 . In fact, multiplying the equation for u_1 (resp. u_{-1}) with u_1 (resp. u_{-1}) and integrating over Ω we obtain

$$\lambda_i(\Omega) = \frac{\int_{\Omega} |\nabla u_i|^2 dx}{\int_{\Omega} u_i^2 dx + \sigma \int_{\partial\Omega} u_i^2 dS} \quad i = -1, 1.$$

Remark 4 Note that from (3.1) we deduce

$$\int_{\Omega} |\nabla u_i|^2 dx = 0$$

and hence $u_i = \text{const.}$, if the denominator

$$\int_{\Omega} u_i^2 dx + \sigma \int_{\partial\Omega} u_i^2 dS = 0.$$

From now on λ and u will denote λ_i and u_i in either case $i = \pm 1$.

Let $(\Omega_t)_t$ be a smooth family of small perturbations of Ω as described in (2.1). In particular they are volume preserving in the sense of (2.4) and (2.7). Denote by u_t the solution to

$$(3.2) \quad \Delta u_t + \lambda(\Omega_t)u_t = 0 \quad \text{in } \Omega_t, \quad \partial_{\nu_t} u_t = \lambda(\Omega_t) \sigma u_t \quad \text{in } \partial\Omega_t.$$

Here $\lambda(\Omega_t)$ has the representation

$$(3.3) \quad \lambda(t) := \lambda(\Omega_t) = \frac{\int_{\Omega_t} |\nabla u_t|^2 dy}{\int_{\Omega_t} u_t^2 dy + \sigma \int_{\partial\Omega_t} u_t^2 dS_t}.$$

As in [5] we transform the integrals onto Ω for small t and differentiate $\lambda(t)$ with respect to t . Then we get

$$\dot{\lambda}(0) = \int_{\partial\Omega} (|\nabla u|^2 - \lambda u^2 - \lambda^2 \sigma^2 u^2 - (n-1)\lambda \sigma H) (v \cdot \nu) dS.$$

In particular $\Omega = B_R$ implies that $\dot{\lambda}(0) = 0$. We are interested in extremality properties of the ball. Thus let $\Omega = B_R$ from now on and let $\sigma < \sigma_0(B_R)$.

We consider the following boundary value problem.

$$(3.4) \quad \Delta u' + \lambda u' = 0 \quad \text{in } B_R \quad \partial_{\nu} u' - \lambda \sigma u' = k(R)(v \cdot \nu) \quad \text{in } \partial B_R$$

where

$$(3.5) \quad k(R) := \lambda u(R) \left(1 + \frac{(n-1)\sigma}{R} + \lambda \sigma^2 \right).$$

We determine the sign of $k(R)$. In this we follow the proof of Lemma 3 in [6]. For the sake of completeness we give the details.

Lemma 2 *Let $k(R)$ be given by (3.5) and let $u(r)$ be the positive radial function in the case $\lambda = \lambda_1$ or $\lambda = \lambda_{-1}$. Then we have*

$$\begin{aligned} k(R) &> 0 && \text{if } \lambda = \lambda_1 \\ k(R) &< 0 && \text{if } \lambda = \lambda_{-1}. \end{aligned}$$

Proof In the radial case either eigenfunction satisfies the differential equation

$$u_{rr} + \frac{n-1}{r} u_r + \lambda u(r) = 0 \quad \text{in } (0, R), \quad u'(R) = \lambda \sigma u(R).$$

We set $z = \frac{u_r}{u}$ and observe that

$$\frac{dz}{dr} + z^2 + \frac{n-1}{r} z + \lambda = 0 \quad \text{in } (0, R).$$

At the endpoint

$$\frac{dz}{dr}(R) + \lambda^2 \sigma^2 + \frac{(n-1)}{R} \lambda \sigma + \lambda = 0.$$

We know that $z(0) = 0$ and $z(R) = \lambda \sigma$. Note that

$$(3.6) \quad z_r(0) = -\lambda.$$

We distinguish two cases.

The case $\lambda = \lambda_1(B_R)$.

In that case we have (also from (3.6))

$$(3.7) \quad z(0) = 0, \quad z(R) = \lambda_1 \sigma < 0, \quad z_r(0) = -\lambda_1 < 0.$$

Thus $z(r)$ decreases near 0. We determine the sign of $z_r(R)$. If $z_r(R) \geq 0$ then because of (3.7) there exists a number $\rho \in (0, R)$ such that $z_r(\rho) = 0$, $z(\rho) < 0$ and $z_{rr}(\rho) \geq 0$. From the equation we get $z_{rr}(\rho) = \frac{n-1}{\rho^2} z(\rho) < 0$ which leads to a contradiction. Consequently

$$z_r(R) = -(\lambda_1^2 \sigma^2 + \frac{(n-1)}{R} \lambda_1 \sigma + \lambda_1) < 0.$$

This implies $k(R) > 0$ in the case $\lambda = \lambda_1(B_R)$.

The case $\lambda = \lambda_{-1}(B_R)$.

In that case we have (see also (3.6))

$$(3.8) \quad z(0) = 0, \quad z(R) = \lambda_{-1} \sigma > 0, \quad z_r(0) = -\lambda_{-1} > 0.$$

Thus $z(r)$ increases near 0. We again determine the sign of $z_r(R)$. If $z_r(R) \leq 0$ then because of (3.8) there exists a number $\rho \in (0, R)$ such that $z_r(\rho) = 0$, $z(\rho) > 0$ and $z_{rr}(\rho) \leq 0$. From the equation we get $z_{rr}(\rho) = \frac{n-1}{\rho^2} z(\rho) > 0$ which is contradictory. Consequently

$$z_r(R) = -(\lambda_{-1}^2 \sigma^2 + \frac{(n-1)}{R} \lambda_{-1} \sigma + \lambda_{-1}) > 0.$$

This also implies $k(R) < 0$ in the case $\lambda = \lambda_{-1}(B_R)$. □

To (3.4) we associate the quadratic form

$$Q(u') := \int_{B_R} |\nabla u'|^2 dx - \lambda \int_{B_R} u'^2 dx - \lambda \sigma \int_{\partial B_R} u'^2 dS.$$

Computations as in [5] lead to the following formula for $\Omega = B_R$. These are the same computations which lead to formula (2.8) in Chapter 2.

$$\ddot{\lambda}(0) = -\lambda^2 \sigma u^2(R) \ddot{S}(0) + \lambda \mathcal{F}$$

where

$$\mathcal{F} = -2Q(u') - 2 \lambda \sigma u(R) k(R) \int_{\partial B_R} (v \cdot \nu)^2 dS.$$

In the remaining part of this chapter we will discuss the sign of $\ddot{\lambda}(0)$.

We modify the approach in Chapter 2.2.1 and consider the following Steklov eigenvalue problem

$$(3.9) \quad \begin{aligned} \Delta \phi + \lambda \phi &= 0 \text{ in } B_R, \\ \partial_\nu \phi - \lambda \sigma \phi &= \mu \phi \text{ on } \partial B_R. \end{aligned}$$

There exists an infinite number of eigenvalues

$$\mu_1 < \mu_2 \leq \mu_3 \leq \dots \lim_{i \rightarrow \infty} \mu_i = \infty.$$

and a complete system of eigenfunctions $\{\phi_i\}_{i \geq 1}$. See also Chapter 7 in [5]. With the same notation we have $u' = \sum_{i=1}^{\infty} c_i \phi_i$ for the solution of (3.4) and we get

$$(3.10) \quad \ddot{\lambda}(0) = \lambda^2 |\sigma| u^2(R) \ddot{S}(0) + 2 \sum_{i=1}^{\infty} c_i^2 \mu_i^2 \left(\frac{\lambda^2 |\sigma| u(R)}{k(R)} - \frac{\lambda}{\mu_i} \right).$$

Let $\lambda = \lambda_1(B_R)$. Then (3.10) is precisely the expression in [5] Chapter 7.2.2 subsection 2. where $\alpha > 0$ is now replaced by $\lambda_1(B_R) |\sigma| > 0$. Thus we conclude

$$(3.11) \quad \ddot{\lambda}_1(0) \geq \lambda_1^2 |\sigma| u^2(R) \ddot{S}(0) > 0.$$

This is a local version of (P5).

The case $\lambda = \lambda_{-1}(B_R) < 0$ is more involved. In that case $\mathcal{F} < 0$ since $k(R) < 0$ by Lemma 2. It is an open problem to show that also in this case $\ddot{\lambda}_{-1}(0) > 0$ - at least for σ close to $\sigma_0(B_R) = -\frac{R}{n}$. This conjecture is motivated by (P6).

4 Harmonic transplantation

The eigenvalues λ_1 (resp. λ_{-1}) have a variational characterization for $\sigma \neq \sigma_0(\Omega)$ (see [4]). Let

$$(4.1) \quad \mathcal{K}_\Omega := \left\{ v \in W^{1,2}(\Omega) : \int_{\Omega} |\nabla v|^2 dx = 1, \int_{\Omega} v dx + \sigma \int_{\partial\Omega} v dS = 0 \right\}.$$

Then for $\sigma < \sigma_0(\Omega)$

$$0 \leq \lambda_1(\Omega) = \frac{1}{\sup \left\{ \int_{\Omega} v^2 dx - |\sigma| \int_{\partial\Omega} v^2 dS : v \in \mathcal{K}_\Omega \right\}}$$

has a unique minimizer (of constant sign). The same holds in the case $0 > \sigma > \sigma_0(\Omega)$ for

$$0 \geq \lambda_{-1}(\Omega) = \frac{1}{\inf \left\{ \int_{\Omega} v^2 dx - |\sigma| \int_{\partial\Omega} v^2 dS : v \in \mathcal{K}_{\Omega} \right\}}.$$

We shortly review the method of harmonic transplantation which has been devised by Hersch[8], (cf. also [2]). In [6] it applied to some shape optimization problems involving Robin eigenvalues. To this end we need the Green's function with Dirichlet boundary condition

$$(4.2) \quad G_{\Omega}(x, y) = \gamma(S(|x - y|) - H(x, y)),$$

where

$$(4.3) \quad \gamma = \begin{cases} \frac{1}{2\pi} & \text{if } n = 2 \\ \frac{1}{(n-2)|\partial B_1|} & \text{if } n > 2 \end{cases} \quad \text{and} \quad S(t) = \begin{cases} -\ln(t) & \text{if } n = 2 \\ t^{2-n} & \text{if } n > 2. \end{cases}$$

For fixed $y \in \Omega$ the function $H(\cdot, y)$ is harmonic.

Definition 1 *The harmonic radius at a point $y \in \Omega$ is given by*

$$r(y) = \begin{cases} e^{-H(y,y)} & \text{if } n = 2, \\ H(y, y)^{-\frac{1}{n-2}} & \text{if } n > 2. \end{cases}$$

The harmonic radius vanishes on the boundary $\partial\Omega$ and takes its maximum r_{Ω} at the harmonic center y_h . It satisfies the isoperimetric inequality [8],[2]

$$(4.4) \quad |B_{r_{\Omega}}| \leq |\Omega|.$$

Note that $G_{B_R}(x, 0)$ is a monotone function in $r = |x|$. Consider any radial function $\phi : B_{r_{\Omega}} \rightarrow \mathbb{R}$ thus $\phi(x) = \phi(r)$. Then there exists a function $\omega : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\phi(x) = \omega(G_{B_{r_{\Omega}}}(x, 0)).$$

To $\phi(x)$ we associate the transplanted function $U : \Omega \rightarrow \mathbb{R}$ defined by $U(x) = \omega(G_{\Omega}(x, y_h))$. Then for any positive function $f(s)$, the following inequalities hold true

$$(4.5) \quad \int_{\Omega} |\nabla U|^2 dx = \int_{B_{r_{\Omega}}} |\nabla \phi|^2 dx$$

$$(4.6) \quad \int_{\Omega} f(U) dx \geq \int_{B_{r_{\Omega}}} f(\phi) dx.$$

$$(4.7) \quad \int_{\Omega} f(U) dx \leq \gamma^n \int_{B_{r_{\Omega}}} f(\phi) dx,$$

where

$$\gamma = \left(\frac{|\Omega|}{|B_{r_{\Omega}}|} \right)^{\frac{1}{n}}.$$

For a proof see [8] or [2] and in particular [6] for a proof of (4.7). The following observation will be useful in the sequel.

Remark 5 Since U is constant on $\partial\Omega$ ($U = U(\partial\Omega)$) and since ϕ is radial we deduce

$$\int_{\partial\Omega} U^2 dS = U^2(\partial\Omega) |\partial\Omega| = \phi^2(r_\Omega) |\partial B_{r_\Omega}| \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} = \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} \int_{\partial B_{r_\Omega}} \phi^2 dS.$$

Let u be a positive radial eigenfunction of $\lambda_1(B_R)$. Then $U \in \mathcal{K}_\Omega$ since $u \in \mathcal{K}_{B_{r_\Omega}}$ and (4.5) holds.

The case $\sigma < \sigma_0 < 0$. By the variational characterization we observe that

$$0 \leq \lambda_1(\Omega) =: \lambda_1^\sigma(\Omega) \leq \frac{1}{\int_{\Omega} U^2 dx - |\sigma| \int_{\partial\Omega} U^2 dS}.$$

We use (4.6) for the first integral in the denominator and Remark 5 for the second.

$$0 \leq \lambda_1^\sigma(\Omega) \leq \frac{1}{\int_{B_{r_\Omega}} u^2 dx - |\sigma| \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} \int_{\partial B_{r_\Omega}} u^2 dS}.$$

Set

$$\sigma' = \sigma \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|}$$

Then we have

$$0 \leq \lambda_1^\sigma(\Omega) \leq \lambda_1^{\sigma'}(B_{r_\Omega}).$$

Alternatively we may write

$$\begin{aligned} & \frac{1}{\int_{B_{r_\Omega}} u^2 dx - |\sigma| \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} \int_{\partial B_{r_\Omega}} u^2 dS} \\ &= \frac{1}{\int_{B_{r_\Omega}} u^2 dx - |\sigma| \int_{\partial B_{r_\Omega}} u^2 dS} \left(1 - \frac{|\sigma| \int_{\partial B_{r_\Omega}} u^2 dS}{\int_{B_{r_\Omega}} u^2 dx - |\sigma| \int_{\partial B_{r_\Omega}} u^2 dS} \left(\frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} - 1 \right) \right)^{-1} \end{aligned}$$

Thus

$$(4.8) \quad 0 \leq \lambda_1^\sigma(\Omega) \leq \lambda_1^{\sigma'}(B_{r_\Omega}) \underbrace{\left(1 - \frac{|\sigma| \int_{\partial B_{r_\Omega}} u^2 dS}{\lambda_1^\sigma(B_{r_\Omega}) \int_{\partial B_{r_\Omega}} u^2 dS} \left(\frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} - 1 \right) \right)^{-1}}_{=: A(\Omega, \sigma)}.$$

Note that the multiplicative term on the right hand side is close to one if the isoperimetric defect $\frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} - 1$ is small. In fact for $\Omega = B_R$ we have $r_\Omega = R$ and thus $A(B_R, \sigma) = 1$.

The case $\sigma_0 < \sigma < 0$. Again by the variational characterization we have

$$0 \geq \lambda_{-1}(\Omega) \geq \frac{1}{\int_{\Omega} U^2 dx - |\sigma| \int_{\partial\Omega} U^2 dS}.$$

We apply (4.7) to the first integral in the denominator and again Remark 5 to the second.

$$\begin{aligned} 0 \geq \lambda_{-1}(\Omega) &\geq \frac{1}{\gamma^n \int_{B_{r_\Omega}} u^2 dx - |\sigma| \frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} \int_{\partial\Omega} u^2 dS} \\ &\geq \frac{1}{\gamma^n \int_{B_{r_\Omega}} u^2 dx - |\sigma| \gamma^n \int_{\partial\Omega} u^2 dS}. \end{aligned}$$

The last inequality holds since the isoperimetric inequality

$$\frac{|\partial\Omega|}{|\partial B_{r_\Omega}|} \geq \frac{|\Omega|}{|B_{r_\Omega}|} = \gamma^n$$

was applied. Thus

$$0 \geq \lambda_{-1}(\Omega) \geq \frac{1}{\gamma^n} \lambda_{-1}(B_{r_\Omega})$$

We may rewrite this as

$$|\Omega| \lambda_{-1}(\Omega) \geq |B_{r_\Omega}| \lambda_{-1}(B_{r_\Omega}).$$

This proves the following theorem.

Theorem 3 *Let Ω be any domain for which the trace operator $W^{1,2}(\Omega) \rightarrow L^2(\partial\Omega)$ is well defined. Let $\lambda_{\pm 1}(\Omega)$ be the first positive (negative) eigenvalue of (3.1) and let r_Ω be the harmonic radius of Ω . Then the following optimality result holds.*

- 1) *In the case $\sigma < \sigma_0(\Omega) < 0$ we have $0 \leq \lambda_1(\Omega) \leq A(\Omega, \sigma) \lambda_1(B_{r_\Omega})$, where the factor A is given in (4.8) and $A = 1$ for the ball.*
- 2) *In the case $\sigma_0(\Omega) < \sigma < 0$ we have $0 \geq |\Omega| \lambda_{-1}(\Omega) \geq |B_{r_\Omega}| \lambda_{-1}(B_{r_\Omega})$.*

Remark 6 *It is interesting to compare 1) in Theorem 3 with (P5). We get the following two sided bounds.*

If B_R is a ball of equal volume with Ω and if $\sigma < \sigma_0(B_R) < 0$ then

$$\lambda_1(B_R) \leq \lambda_1(\Omega) \leq A(\Omega, \sigma) \lambda_1(B_{r_\Omega}).$$

Equality holds for the ball.

Acknowledgement This paper was written during a visit at the Newton Institute in Cambridge. Both authors would like to thank this Institute for the excellent working atmosphere.

References

- [1] G. Auchmuty, *Sharp boundary trace inequalities*, Proc. Royal Soc. Edinburgh A, 144 (2014), 1-12.
- [2] C. Bandle and M. Flucher, *Harmonic radius and concentration of energy; hyperbolic radius and Liouville's equations $\Delta U = e^U$ and $\Delta U = U^{(n+2)/(n-2)}$* , SIAM Review 38 (1996), 191-238.
- [3] C. Bandle, *A Rayleigh-Faber-Krahn inequality and some monotonicity properties for eigenvalue problems with mixed boundary conditions*, Birkhäuser ISNM 157 (2008), 3-12.
- [4] C. Bandle, J. v. Below and W. Reichel, *Parabolic problems with dynamical boundary conditions: eigenvalue expansions and blow up*. Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 17 (2006), no. 1, 3567.
- [5] C. Bandle and A. Wagner, *Second variation of domain functionals and applications to problems with Robin boundary conditions*, arXiv (2014)
- [6] C. Bandle and A. Wagner, *Isoperimetric inequalities for the principal eigenvalue of a membrane and the energy of problems with Robin boundary conditions*, to appear in Journal of Convex Analysis.
- [7] D. Bucur and A. Giacomini, *Faber-Krahn inequalities for the Robin-Laplacian: a free discontinuity approach*,
- [8] J. Hersch, *Transplantation harmonique, transplantation par modules, et théorèmes isopérimétriques*, Comment. Math. Helv. 44 (1969), 354-366.

Reports des Instituts für Mathematik der RWTH Aachen

- [1] Bemelmans J.: *Die Vorlesung "Figur und Rotation der Himmelskörper" von F. Hausdorff, WS 1895/96, Universität Leipzig*, S 20, März 2005
- [2] Wagner A.: *Optimal Shape Problems for Eigenvalues*, S 30, März 2005
- [3] Hildebrandt S. and von der Mosel H.: *Conformal representation of surfaces, and Plateau's problem for Cartan functionals*, S 43, Juli 2005
- [4] Reiter P.: *All curves in a C^1 -neighbourhood of a given embedded curve are isotopic*, S 8, Oktober 2005
- [5] Maier-Paape S., Mischaikow K. and Wanner T.: *Structure of the Attractor of the Cahn-Hilliard Equation*, S 68, Oktober 2005
- [6] Strzelecki P. and von der Mosel H.: *On rectifiable curves with L^p bounds on global curvature: Self-avoidance, regularity, and minimizing knots*, S 35, Dezember 2005
- [7] Bandle C. and Wagner A.: *Optimization problems for weighted Sobolev constants*, S 23, Dezember 2005
- [8] Bandle C. and Wagner A.: *Sobolev Constants in Disconnected Domains*, S 9, Januar 2006
- [9] McKenna P.J. and Reichel W.: *A priori bounds for semilinear equations and a new class of critical exponents for Lipschitz domains*, S 25, Mai 2006
- [10] Bandle C., Below J. v. and Reichel W.: *Positivity and anti-maximum principles for elliptic operators with mixed boundary conditions*, S 32, Mai 2006
- [11] Kyed M.: *Travelling Wave Solutions of the Heat Equation in Three Dimensional Cylinders with Non-Linear Dissipation on the Boundary*, S 24, Juli 2006
- [12] Blatt S. and Reiter P.: *Does Finite Knot Energy Lead To Differentiability?*, S 30, September 2006
- [13] Grunau H.-C., Ould Ahmedou M. and Reichel W.: *The Paneitz equation in hyperbolic space*, S 22, September 2006
- [14] Maier-Paape S., Miller U., Mischaikow K. and Wanner T.: *Rigorous Numerics for the Cahn-Hilliard Equation on the Unit Square*, S 67, Oktober 2006
- [15] von der Mosel H. and Winklmann S.: *On weakly harmonic maps from Finsler to Riemannian manifolds*, S 43, November 2006
- [16] Hildebrandt S., Maddocks J. H. and von der Mosel H.: *Obstacle problems for elastic rods*, S 21, Januar 2007
- [17] Galdi P. Giovanni: *Some Mathematical Properties of the Steady-State Navier-Stokes Problem Past a Three-Dimensional Obstacle*, S 86, Mai 2007
- [18] Winter N.: *$W^{2,p}$ and $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear, uniformly elliptic equations*, S 34, Juli 2007
- [19] Strzelecki P., Szumańska M. and von der Mosel H.: *A geometric curvature double integral of Menger type for space curves*, S 20, September 2007
- [20] Bandle C. and Wagner A.: *Optimization problems for an energy functional with mass constraint revisited*, S 20, März 2008
- [21] Reiter P., Felix D., von der Mosel H. and Alt W.: *Energetics and dynamics of global integrals modeling interaction between stiff filaments*, S 38, April 2008
- [22] Belloni M. and Wagner A.: *The ∞ Eigenvalue Problem from a Variational Point of View*, S 18, Mai 2008
- [23] Galdi P. Giovanni and Kyed M.: *Steady Flow of a Navier-Stokes Liquid Past an Elastic Body*, S 28, Mai 2008
- [24] Hildebrandt S. and von der Mosel H.: *Conformal mapping of multiply connected Riemann domains by a variational approach*, S 50, Juli 2008
- [25] Blatt S.: *On the Blow-Up Limit for the Radially Symmetric Willmore Flow*, S 23, Juli 2008
- [26] Müller F. and Schikorra A.: *Boundary regularity via Uhlenbeck-Rivière decomposition*, S 20, Juli 2008
- [27] Blatt S.: *A Lower Bound for the Gromov Distortion of Knotted Submanifolds*, S 26, August 2008
- [28] Blatt S.: *Chord-Arc Constants for Submanifolds of Arbitrary Codimension*, S 35, November 2008
- [29] Strzelecki P., Szumańska M. and von der Mosel H.: *Regularizing and self-avoidance effects of integral Menger curvature*, S 33, November 2008
- [30] Gerlach H. and von der Mosel H.: *Yin-Yang-Kurven lösen ein Packungsproblem*, S 4, Dezember 2008
- [31] Buttazzo G. and Wagner A.: *On some Rescaled Shape Optimization Problems*, S 17, März 2009
- [32] Gerlach H. and von der Mosel H.: *What are the longest ropes on the unit sphere?*, S 50, März 2009
- [33] Schikorra A.: *A Remark on Gauge Transformations and the Moving Frame Method*, S 17, Juni 2009
- [34] Blatt S.: *Note on Continuously Differentiable Isotopies*, S 18, August 2009
- [35] Knappmann K.: *Die zweite Gebietsvariation für die gebeulte Platte*, S 29, Oktober 2009
- [36] Strzelecki P. and von der Mosel H.: *Integral Menger curvature for surfaces*, S 64, November 2009
- [37] Maier-Paape S., Imkeller P.: *Investor Psychology Models*, S 30, November 2009
- [38] Scholtes S.: *Elastic Catenoids*, S 23, Dezember 2009
- [39] Bemelmans J., Galdi G.P. and Kyed M.: *On the Steady Motion of an Elastic Body Moving Freely in a Navier-Stokes Liquid under the Action of a Constant Body Force*, S 67, Dezember 2009
- [40] Galdi G.P. and Kyed M.: *Steady-State Navier-Stokes Flows Past a Rotating Body: Leray Solutions are Physically Reasonable*, S 25, Dezember 2009

- [41] Galdi G.P. and Kyed M.: *Steady-State Navier-Stokes Flows Around a Rotating Body: Leray Solutions are Physically Reasonable*, S 15, Dezember 2009
- [42] Bemelmans J., Galdi G.P. and Kyed M.: *Fluid Flows Around Floating Bodies, I: The Hydrostatic Case*, S 19, Dezember 2009
- [43] Schikorra A.: *Regularity of $n/2$ -harmonic maps into spheres*, S 91, März 2010
- [44] Gerlach H. and von der Mosel H.: *On sphere-filling ropes*, S 15, März 2010
- [45] Strzelecki P. and von der Mosel H.: *Tangent-point self-avoidance energies for curves*, S 23, Juni 2010
- [46] Schikorra A.: *Regularity of $n/2$ -harmonic maps into spheres (short)*, S 36, Juni 2010
- [47] Schikorra A.: *A Note on Regularity for the n -dimensional H -System assuming logarithmic higher Integrability*, S 30, Dezember 2010
- [48] Bemelmans J.: *Über die Integration der Parabel, die Entdeckung der Kegelschnitte und die Parabel als literarische Figur*, S 14, Januar 2011
- [49] Strzelecki P. and von der Mosel H.: *Tangent-point repulsive potentials for a class of non-smooth m -dimensional sets in \mathbb{R}^n . Part I: Smoothing and self-avoidance effects*, S 47, Februar 2011
- [50] Scholtes S.: *For which positive p is the integral Menger curvature \mathcal{M}_p finite for all simple polygons*, S 9, November 2011
- [51] Bemelmans J., Galdi G. P. and Kyed M.: *Fluid Flows Around Rigid Bodies, I: The Hydrostatic Case*, S 32, Dezember 2011
- [52] Scholtes S.: *Tangency properties of sets with finite geometric curvature energies*, S 39, Februar 2012
- [53] Scholtes S.: *A characterisation of inner product spaces by the maximal circumradius of spheres*, S 8, Februar 2012
- [54] Kolasiński S., Strzelecki P. and von der Mosel H.: *Characterizing $W^{2,p}$ submanifolds by p -integrability of global curvatures*, S 44, März 2012
- [55] Bemelmans J., Galdi G.P. and Kyed M.: *On the Steady Motion of a Coupled System Solid-Liquid*, S 95, April 2012
- [56] Deipenbrock M.: *On the existence of a drag minimizing shape in an incompressible fluid*, S 23, Mai 2012
- [57] Strzelecki P., Szumańska M. and von der Mosel H.: *On some knot energies involving Menger curvature*, S 30, September 2012
- [58] Overath P. and von der Mosel H.: *Plateau's problem in Finsler 3-space*, S 42, September 2012
- [59] Strzelecki P. and von der Mosel H.: *Menger curvature as a knot energy*, S 41, Januar 2013
- [60] Strzelecki P. and von der Mosel H.: *How averaged Menger curvatures control regularity and topology of curves and surfaces*, S 13, Februar 2013
- [61] Hafizogullari Y., Maier-Paape S. and Platen A.: *Empirical Study of the 1-2-3 Trend Indicator*, S 25, April 2013
- [62] Scholtes S.: *On hypersurfaces of positive reach, alternating Steiner formulæ and Hadwiger's Problem*, S 22, April 2013
- [63] Bemelmans J., Galdi G.P. and Kyed M.: *Capillary surfaces and floating bodies*, S 16, Mai 2013
- [64] Bandle, C. and Wagner A.: *Domain derivatives for energy functionals with boundary integrals; optimality and monotonicity.*, S 13, Mai 2013
- [65] Bandle, C. and Wagner A.: *Second variation of domain functionals and applications to problems with Robin boundary conditions*, S 33, Mai 2013
- [66] Maier-Paape, S.: *Optimal f and diversification*, S 7, Oktober 2013
- [67] Maier-Paape, S.: *Existence theorems for optimal fractional trading*, S 9, Oktober 2013
- [68] Scholtes, S.: *Discrete Möbius Energy*, S 11, November 2013
- [69] Bemelmans, J.: *Optimale Kurven – über die Anfänge der Variationsrechnung*, S 22, Dezember 2013
- [70] Scholtes, S.: *Discrete Thickness*, S 12, Februar 2014
- [71] Bandle, C. and Wagner A.: *Isoperimetric inequalities for the principal eigenvalue of a membrane and the energy of problems with Robin boundary conditions.*, S 12, März 2014
- [72] Overath P. and von der Mosel H.: *On minimal immersions in Finsler space.*, S 26, April 2014
- [73] Bandle, C. and Wagner A.: *Two Robin boundary value problems with opposite sign.*, S 17, Juni 2014